

A leptin fragment mirrors the cognitive enhancing and neuroprotective actions of leptin.

Yasaman Malekizadeh¹, Alison Holiday², Devon Redfearn², James A Ainge², Gayle Doherty², Jenni Harvey¹

¹ Division of Neuroscience, Medical Research Institute, Ninewells Hospital and Medical School, University of Dundee, Dundee DD1 9SY; ²School of Psychology and Neuroscience, St Mary's Quad, University of St Andrews, St Andrews, Fife, KY16 9TS

Corresponding author: Dr Jenni Harvey, Division of Neuroscience, Medical Research Institute, Ninewells Hospital and Medical School, University of Dundee, Dundee DD1 9SY

Tel: +44 1382 383359

Email: j.z.harvey@dundee.ac.uk

Running Title: Leptin (116-130) has cognitive enhancing and neuroprotective actions

Keywords: Hippocampus, synaptic plasticity, amyloid beta, AMPA receptor trafficking, episodic memory, Alzheimer's disease

Abstract:

A key pathology of Alzheimer's disease (AD) is amyloid β ($A\beta$) accumulation which triggers synaptic impairments and neuronal death. Metabolic disruption is common in AD and recent evidence implicates impaired leptin function in AD. Thus the leptin system may be a novel therapeutic target in AD. Indeed, leptin has cognitive enhancing properties and it prevents the aberrant effects of $A\beta$ on hippocampal synaptic function and neuronal viability. However as leptin is a large peptide, development of smaller leptin-mimetics may be the best therapeutic approach. Thus, we have examined the cognitive enhancing and neuroprotective properties of known bioactive leptin fragments. Here we show that the leptin (116-130) fragment, but not leptin (22-56), mirrored the ability of leptin to promote AMPA receptor trafficking to synapses and facilitate activity-dependent hippocampal synaptic plasticity. Administration of leptin (116-130) also mirrored the cognitive enhancing effects of leptin as it enhanced performance in episodic-like memory tests. Moreover, leptin (116-130) prevented hippocampal synaptic disruption and neuronal cell death in models of amyloid toxicity. These findings establish further the importance of the leptin system as a therapeutic target in AD.

Introduction.

Alzheimer's disease (AD) is a complex, progressive brain disorder that results in profound cognitive deficits, particularly in memory. Accumulation of toxic amyloid plaques and neurofibrillary tangles comprising hyper-phosphorylated tau are key pathological features of AD. Proteolytic processing of APP and generation of toxic amyloid beta ($A\beta$) is linked to aberrant synaptic function and neuronal degeneration. Several studies indicate that soluble $A\beta$ oligomers greatly influence activity-dependent hippocampal synaptic plasticity resulting in deficits in learning and memory. Indeed, $A\beta$ inhibits the induction of hippocampal long-term

potentiation (LTP), but facilitates long-term depression (LTD; Shankar et al, 2008; Walsh et al, 2002). Furthermore, A β significantly alters AMPA receptor trafficking to and away from synapses which contributes to hippocampal synaptic dysfunction (Hsieh et al, 2006; Liu et al, 2010).

Clinical evidence indicates that diet and lifestyle are major risk factors for developing AD and disruption to metabolic systems is linked to AD (Stranahan and Mattson, 2012). The metabolic hormone leptin regulates energy homeostasis, but also markedly influences hippocampal synaptic function. Indeed, leptin-insensitive rodents exhibit impaired hippocampal LTP and spatial memory (Li et al, 2002). Moreover, direct administration of leptin into the hippocampus facilitates learning (Wayner et al, 2004), whereas hippocampal dendritic morphology, glutamate receptor trafficking and synaptic plasticity are significantly altered after leptin treatment (Shanley et al, 2001; O'Malley et al, 2007; Moulton et al, 2010; Luo et al, 2015). Clinical studies indicate that aberrant leptin function is associated with an increased risk of AD, as AD patients exhibit significantly lower leptin levels than normal (Power et al, 2001). Individuals with lower circulating leptin also have a greater risk of developing AD (Lieb et al, 2009). Additionally, leptin levels are markedly reduced in rodent models with familial AD mutations (APP^{Swe}; PS1^{M146V}; Fewlass et al, 2004). In cellular models of AD, leptin treatment prevents the aberrant effects of A β , as leptin reverses the ability of A β to inhibit LTP and facilitate LTD in hippocampal slices (Doherty et al, 2013) and it prevents A β -driven internalisation of the AMPA receptor subunit, GluA1 in hippocampal neurons (Doherty et al, 2013). Chronic intracerebroventricular injection of leptin also reverses A β -induced impairments in LTP *in vivo* (Tong et al, 2015). In cell survival assays, leptin protects against A β -induced toxicity as neuronal viability is increased after leptin treatment. Moreover the cortical expression of two AD-linked biomarkers, endophilin 1 and phosphorylated tau, are enhanced in leptin-insensitive Zucker *fa/fa* rats

(Doherty et al, 2013). Thus there is growing evidence that leptin-based therapies may be beneficial in AD.

However, using leptin therapeutically may not be the best approach due to its widespread central actions. One possibility is to develop small molecules that mimic leptin action.

Indeed, studies have found that specific fragments of the leptin peptide are bioactive and mirror the anti-obesity effects of leptin (Grasso et al, 1997; Roshavskaya-Arena et al, 2000).

Indeed, application of a C-terminal fragment of leptin (amino acids 116-130) or a shorter fragment (116-121) to leptin-deficient *ob/ob* mice reduced food intake and body weight

(Grasso et al, 1997; Roshavskaya-Arena et al, 2000; Grasso et al, 2001). However, another leptin fragment (22-56) is also bioactive as significant reductions in food intake have been

observed after leptin (22-56) administration (Samson et al, 1996). Thus, it is feasible that

different parts of the leptin molecule are active in the CNS. However, the cognitive

enhancing and neuroprotective properties of the bioactive leptin fragments are unknown.

Here we show that leptin (116-130), but not leptin (22-56), mirrored the effects of leptin on hippocampal synaptic plasticity and also prevented hippocampal synaptic disruption and cell

death induced by A β . Moreover, leptin (116-130) has similar cognitive enhancing effects to

leptin as it enhanced episodic-like memory. These findings establish further the importance

of the leptin system as a therapeutic target in AD.

Methods.

Primary Neuronal culture.

Hippocampal cultures were prepared from neonatal Sprague Dawley rats as before (O'Malley et al, 2007). Briefly, neonatal Sprague Dawley rats (1–3 days old) were killed by cervical dislocation in accordance with Schedule 1 of the United Kingdom Government Animals

(Scientific Procedures) Act, 1986 and the hippocampi removed. After washing in HEPES buffered saline (HBS) comprising (mM): NaCl 135; KCl 5; CaCl₂ 1; MgCl₂ 1; HEPES 10; D-glucose 25 at pH 7.4; cells were treated with protease Type X and Type XIV (0.5 mg/ml; Sigma) for 25 min at room temperature. Dissociated cells were plated onto sterile dishes (Falcon 3001) treated with poly-L-lysine (20 µg/ml; 1-2 h) and maintained in MEM with serum replacement-2 (Sigma) in a humidified atmosphere of 5% CO₂ and 95% O₂ at 37 °C for up to 3 weeks.

Human Neural Cell Line SH-SY5Y

The human neuroblastoma cell line, SH-SY5Y (ECACC, UK) was maintained in Dulbecco's Modified Eagle Medium supplemented with glucose (4500µg/l) and 10% (v/v) cosmic calf serum (Fisher Scientific, UK) at 37°C in a humidified atmosphere of 5% CO₂, 95% air and allowed to reach 70-80% confluence before seeding. Cells (passage 10-18) were plated at 10,000 cells per well in 96 well tissue culture plates (Nunc, VWR, UK) and at a density of 2X10⁵ cells in 35mm dishes for protein extraction. To induce differentiation, cells were cultured in DMEM supplemented with glucose (4500µg/l), 1% (v/v) cosmic calf serum and 10µM retinoic acid for 5 days. Thereafter they were incubated in DMEM supplemented with glucose (4500µg/l), serum replacement 2 (2%; Sigma, UK) and 18µM 5-fluorodeoxyuridine to inhibit proliferation of undifferentiated cells. 50% of the medium was changed every 2-3 days and pharmacological treatment was carried out 7 days after switching to this medium. Reagents used were (from Sigma UK unless stated) 0.1-10nM human leptin and leptin (116-130); 0.1-10nM leptin (22-56; Bachem; Switzerland); 5µM copper chloride; 10mM Aβ₁₋₄₂; 5µM WP1066 or 50nM wortmannin. All treatments were added to the culture at the same time and survival assays were carried out after 96 hours treatment. Protein samples for ELISA were extracted 3 hours after exposure to the relevant reagents

Cell Survival Assays

The concentration of lactate dehydrogenase in the culture medium (LDH) was used to monitor the level of cell death, as previously (Oldrieve and Doherty, 2010). A Crystal violet assay was used to assess total cell number. Cells were fixed in neutral buffered formalin and washed 3 times in PBS prior to staining with 0.01% crystal violet acetate for 5 min. Plates were washed 5-10 times in dH₂O and cells solubilised in 100µl DMSO before reading the absorbance on a Biohit BP100 plate reader. For both assays, data is expressed as percentage relative to control, untreated wells to normalise for differences in plating density between individual experiments.

ELISA for cell signalling pathways

Protein from cultures was extracted into 500µl Tris-buffered saline containing protease inhibitor cocktail, and a Bradford assay used to determine protein concentration. Samples were diluted to give equal loading onto ELISA plates. Commercially available ELISA kits were used in accordance with the manufacturer's instructions to determine the ratio of pan-STAT3 to phospho-STAT3 (Sigma, UK) and pan-Akt to phospho-Akt (Sigma, UK). Each protein sample was run in duplicate.

Surface labelling of AMPA receptors.

To monitor GluA1 surface expression, immunocytochemistry was performed on hippocampal cultures (7-14 DIV) as before (Moult et al, 2010). Neurons were treated with agents for 20 min at room temperature (20-22 °C) before incubation with an antibody against an N-terminal region of GluA1 (sheep anti-GluR1; in house antibody against synthetic peptide (RTSDSRDHTRVDWKR) corresponding to 253–267 residues of GluR1; 1:100; Moult et al,

2010) at 4°C. Neurons were then fixed with 4% paraformaldehyde (5 min) before adding an appropriate fluorescently-labelled secondary antibody (1:250; 30 min) to enable visualisation of GluA1 surface staining. No labeling was observed after incubation with secondary antibody alone. For synaptic co-localization studies, surface GluA1 staining was compared with synaptophysin (anti-synaptophysin antibody) immunolabelling and percentage of surface GluA1 at synapses was assessed as the number of GluA1-positive sites that colocalize with synaptophysin-positive sites.

An Axiovert 200M laser scanning confocal microscope was used for image acquisition and analysis. Dual labeling images were obtained in multi-tracking mode using a 15 s scan speed. Intensity of staining was determined off-line using Lasersharp software (Carl Zeiss). Analysis lines (50 µm) were drawn along randomly selected dendritic regions (50 nm), and mean fluorescence intensity was calculated. Data were obtained from at least four randomly selected cells for each condition, and all data were obtained from at least three different cultures from different animals. Within a given experiment, all conditions, including illumination intensity and photomultiplier gains, were kept constant. To quantify experimental data obtained from separate days, data were normalized relative to mean fluorescence intensity in control neurons.

Hippocampal slice preparation and electrophysiology.

Parasagittal hippocampal slices (300 µm) were prepared from either P13-21 or 12-24 week old male Sprague Dawley rats as previously (Moult et al, 2010). Brains were rapidly removed and placed in ice-cold artificial CSF (aCSF; bubbled with 95% O₂ and 5% CO₂) containing the following (in mM): 124 NaCl, 3 KCl, 26 NaHCO₃, 1.25 NaH₂PO₄, 2 CaCl₂, 1 MgSO₄, and 10 d-glucose. Once prepared, parasagittal slices were allowed to recover at room temperature in oxygenated aCSF for 1 h before use. Slices were transferred to a submerged

chamber maintained at room temperature and perfused with artificial cerebrospinal fluid at 2mlmin^{-1} . Standard extracellular recordings were used to monitor evoked field excitatory postsynaptic potentials (fEPSP) from stratum radiatum. The Schaffer collateral-commissural pathway was stimulated (constant voltage; 0.1ms) at 0.033 Hz , using a stimulus intensity that evoked a peak amplitude $\sim 50\%$ of the maximum. Synaptic potentials were low pass filtered at 2 kHz and digitally sampled at 10 kHz . The fEPSP slope was measured and expressed relative to baseline. Synaptic records are the average of four consecutive responses and stimulus artefacts are blanked for clarity.

LTP was induced using a high frequency stimulation paradigm (10 trains; 8 stimuli; 100Hz), whereas the sub-threshold LTD paradigm used was low frequency stimulation (LFS; 1 Hz) for 5 min (300 shocks; 15). A primed burst stimulation paradigm (5 trains of 8 stimuli at 100Hz ; Rose and Dunwiddie, 1986) was used to induce STP. Recordings were made using an Axopatch 200B amplifier and analysed using LTP v2.4 software (Anderson and Collingridge, 2007). In all experiments, the slope of fEPSPs was monitored continuously. For studies comparing the actions of leptin and the fragments on synaptic transmission, the mean slope (average of 5 min recording) of fEPSPs obtained during the 5 min period immediately prior to leptin and/or agent addition was compared to that after $25\text{-}30\text{ min}$ exposure to the agent. In synaptic plasticity studies, the degree of potentiation or depression was calculated $30\text{-}35\text{ min}$ after LFS or HFS and expressed as a percentage of baseline $\pm\text{ SEM}$.

Episodic-like memory testing

Mice were tested using the same protocol used to test object-place-context (OPC) recognition in rats (Wilson et al, 2013). The size of the testing box was reduced to $30\text{cm} \times 20\text{cm} \times 20\text{cm}$ and the household objects used in the task were approximately the size of the mice in at least one dimension. The box could be configured with two sets of contextual features (white vs.

stripes) by using different wall and floor inserts. These created one environment that was plain white and one that had stripes and a mesh floor. Mice were handled extensively prior to training and then habituated to the box for 4 days before undergoing object recognition, object-place and object-context training. These data are not presented and served only to familiarise the mice to the testing procedures and the factors that are manipulated during the OPC task. On the OPC testing days, mice were given IP injections of either saline, leptin or leptin (116-130, both at 7.8nM/ml) 30 minutes prior to testing. Testing consisted of three stages; 2 sample phases and a test phase (illustrated in Fig 7A). Each phase was three min duration and exploration behaviour was scored using custom written software (Observe, University of St Andrews). Between all phases the box was cleaned and dried using F10 solution. All testing was recorded on a webcam connected to a laptop to allow offline scoring of behaviour. Twenty percent of trials were rescored by a researcher who was blind to the group identity of the mice and these scores were found to be within 10% of the original scoring. The dependent variable measured was a discrimination index which is calculated by subtracting the amount of time exploring the familiar object from the amount of time exploring the novel object and dividing by the total amount of exploration. This measure was taken from the first minute of the test trial as novelty discrimination has been shown to rapidly diminish after the first minute (Clarke et al., 2000).

Statistical Analyses.

In immunocytochemical studies, data are expressed as mean \pm SEM and statistical analyses performed using one-way ANOVA with Tukey's post hoc test for comparisons between multiple groups. In electrophysiological studies, analyses were performed using repeated measures ANOVA for comparison between multiple groups. A one-way ANOVA with Tukey's post hoc test was used for comparison between multiple groups in cell viability

assays. In ELISA assays, a t-test was used to determine differences between groups. $P < 0.05$ was considered significant. In behavioural studies a one-way ANOVA with Tukey's post hoc tests was used for comparison between the groups.

Materials.

Recombinant human leptin (R&D Systems, Europe) was used in all experiments. Leptin was prepared as a 50 μ M stock solution in normal aCSF and was diluted to the required concentration in aCSF or HBS. A β_{1-42} peptide was synthesized and purified by Dr. Elliott at Yale University (New Haven, CT, USA) based on the human A β sequence. The lyophilized powder was solubilized in dimethyl sulfoxide (DMSO) and diluted to a working concentration. For cell viability and biochemical assays A β was prepared as described previously as A β has shown to elicit toxicity when in a β sheet conformation (Simmons et al., 1994). Briefly, lyophilized powder was solubilized in dH₂O to a concentration of 1 mM in phosphate-buffered saline and incubated for 24 hours at 37 °C prior to further dilution to working concentration in culture medium. Wortmannin, bisperoxovanadium (Calbiochem), WP1066 (Cayman Chemicals) and D-AP5 (Tocris Cookson) were all obtained commercially.

Results

Leptin (116-130) facilitates NMDA receptor-dependent hippocampal synaptic plasticity.

It is known that NMDA receptors contribute little to basal excitatory synaptic transmission but synaptic activation of NMDA receptors is crucial for LTP induction at hippocampal synapses (Bliss and Collingridge, 1993). Previous studies indicate that leptin enhances the magnitude of activity-dependent LTP in acute hippocampal slices (Oomura et al, 2006) and following direct administration into the hippocampus *in vivo* (Wayner et al, 2004). We have

also shown that leptin facilitates NMDA receptor-dependent synaptic plasticity as leptin promotes conversion of short term potentiation (STP) to hippocampal LTP (Shanley et al, 2001). In order to compare the effects of leptin and the leptin fragments on synaptic plasticity in juvenile hippocampal slices (P13-21), a primed burst stimulation paradigm (5 trains of 8 stimuli at 100Hz; Rose and Dunwiddie, 1986) was used to induce STP, which returned to baseline levels within 30-35 min (n=4; Fig 1A). In accordance with previous studies (Shanley et al, 2001), application of leptin (50nM) prior to the stimulation paradigm converted STP into a persistent increase in synaptic transmission ($155 \pm 6.1\%$ of baseline at 40 min; n=4; $p < 0.05$; Fig 1A). Similarly, treatment with 50nM leptin (116-130) facilitated synaptic plasticity as synaptic transmission was enhanced to $143 \pm 6.4\%$ of baseline in leptin (116-130)-treated slices (n=4; $p < 0.05$; Fig 1B). In contrast, in slices exposed to 50nM leptin (22-56), only STP was observed as synaptic transmission returned to baseline levels within 40 min (n=6; Fig 1C).

Leptin (116-130) induces synaptic plasticity at adult hippocampal CA1 synapses.

We have shown that leptin regulation of excitatory synaptic transmission is age-dependent. Thus in contrast to its effects in juvenile tissue, leptin (25nM) induces a novel form of LTP in adult hippocampus (Moult and Harvey, 2011). In order to verify if the leptin fragments mirror leptin action in adult tissue, the effects of the leptin fragments were examined in hippocampal slices from adult (12-24 week) rats. In accordance with previous studies (Moult et al, 2010; Moult and Harvey, 2011), application of leptin (25nM; 20 min) to adult slices rapidly enhanced synaptic transmission (to $188 \pm 13\%$ of baseline; n=4; $p < 0.01$; data not shown) which was sustained for the duration of recordings. Synaptic transmission was also markedly increased (to $140 \pm 13\%$ of baseline; n=5; $p < 0.05$; Fig 2A) after treatment with 25nM leptin (116-130), whereas application of 25nM leptin (22-56) had no significant effect on synaptic efficacy ($99 \pm 2\%$ of baseline; n=4; $p > 0.05$; Fig 2B).

These data indicate that leptin (116-130) mirrors the ability of leptin to enhance synaptic efficacy in adult hippocampus.

Leptin (116-130), but not leptin (22-56) enhances the surface expression of GluA1.

Trafficking of AMPA receptors to and away from synapses is crucial for various forms of activity-dependent synaptic plasticity (Collingridge et al, 2004). Our studies indicate that leptin regulates AMPA receptor trafficking as leptin promotes trafficking of GluA1 to hippocampal synapses (Moult et al, 2010). Moreover the ability of leptin to induce LTP in adult hippocampus requires the delivery of GluA1 to synapses by leptin (Moult et al, 2010). Thus, to assess if the leptin fragments also influence AMPA receptor trafficking processes, the surface expression of GluA1 was assayed in cultured hippocampal neurons (Moult et al, 2010). In agreement with previous studies, application of leptin (50nM; 15min) increased surface GluA1 expression to $184 \pm 7\%$; (n=36; $p < 0.001$; Fig 2C&D) compared to control hippocampal neurons. Similarly GluA1 surface expression was elevated (to $160 \pm 6\%$ of control; n=36; $p < 0.001$) after exposure to 50nM leptin (116-130; Fig 2C&D), whereas GluA1 surface expression ($98 \pm 4\%$ of control; n=36; $p > 0.05$) was not altered after treatment with 50nM leptin (22-56). These data indicate that leptin (116-130), but not leptin (22-56) mirrors the plasticity enhancing effects of leptin by increasing GluA1 surface expression in hippocampal neurons.

As excitatory synaptic strength is governed by the density of AMPA receptors expressed at synapses, the effects on synaptic AMPA receptors was examined by comparing the co-localisation between surface GluA1 and synaptophysin immunolabelling in hippocampal cultures (Moult et al, 2010). In agreement with previous studies (O'Malley et al, 2007; Moult et al, 2010), leptin (50nM; 30 min) increased synaptophysin staining to $144 \pm 9\%$ of control (n=48; $p < 0.001$) and it also enhanced the degree of co-localisation between surface GluA1 and synaptophysin immunostaining

from $43 \pm 5.4\%$ to $62 \pm 4.4\%$ ($n=36$; $p<0.05$; Fig 2E). Similarly, in hippocampal neurons treated with leptin (116-130), synaptophysin staining was increased to $122 \pm 9\%$ ($n=36$; $p<0.05$) and a significant increase (to $64 \pm 4.9\%$; $n=36$; $p<0.01$) in GluA1-synaptophysin co-localisation was observed (Fig 2E). Conversely, exposure to leptin (22-56) had no significant effect on either synaptophysin staining ($105 \pm 9\%$; $n=36$; $p>0.05$) or the degree of co-localisation ($49 \pm 5.4\%$; $n=36$; $p>0.05$; Fig 2E). These data indicate that leptin (116-130), but not leptin (22-56) promotes the trafficking of GluA1 to synapses in hippocampal neurons.

Inhibition of the phosphatase, PTEN underlies leptin-driven trafficking of GluA1 to synapses (Moult et al, 2010). In order to determine if similar leptin dependent signalling cascades mediate the actions of leptin (116-130), the effects of pharmacological inhibition of PTEN with bisperoxovanadium (bpV; Schmid et al, 2004) were assessed in hippocampal cultures. Application of bpV (50nM; 30 min) increased GluA1 surface expression to $148 \pm 8\%$ of control ($n=36$; $p<0.001$; Fig 2F). In accordance with previous studies (Moult et al, 2010), leptin resulted in a significant increase in surface GluA1 labelling ($140 \pm 7.2\%$; $n=36$; $p<0.001$; Fig 2F). Similarly, GluA1 surface expression was enhanced (to $167 \pm 10\%$ of control; $n=36$; $p<0.001$) after exposure to leptin (116-130). Moreover, treatment with bpV occluded the effects of either leptin or leptin (116-130) such that GluA1 staining was $131 \pm 6.6\%$ and $148 \pm 8.1\%$ of control ($n=36$; $p<0.001$; for both; Fig 2F) in the presence of bpV and either leptin or leptin (116-130), respectively. These data indicate that the ability of leptin (116-130) to regulate GluA1 trafficking involves inhibition of PTEN.

Leptin (116-130), but not leptin (22-56) reverses $A\beta_{1-42}$ inhibition of hippocampal synaptic plasticity

Several studies indicate that soluble A β oligomers impair activity-dependent synaptic plasticity, as exposure to A β inhibits hippocampal LTP (Shankar et al, 2008; Li et al, 2009) and enhances LTD (Shankar et al, 2008). Moreover, our recent studies indicate that leptin reverses the detrimental effects of A β_{1-42} on both LTP and LTD (Doherty et al, 2013). Thus we assessed if either of the fragments mirrored the protective actions of leptin on hippocampal synaptic plasticity. Initially we determined that application of leptin prevented the acute effects of A β_{1-42} on LTP. In control slices, synaptic plasticity was induced using high frequency stimulation (HFS; 100 Hz 10 trains of 8 stimuli) which increased synaptic transmission to $127 \pm 3.4\%$ of baseline (n=8; p<0.01). Similarly in slices treated with the inactive peptide A β_{42-1} (1 μ M; 40 min), an enhancement of synaptic transmission ($132 \pm 8.8\%$ of baseline) was induced (n=5; p<0.05; Figure 3A). However in accordance with previous studies (Doherty et al, 2013), exposure to A β_{1-42} not only blocked hippocampal synaptic plasticity ($96 \pm 7.5\%$ of baseline; n=5; p>0.05; Fig 3A) but this effect was reversed by leptin (25nM; $135 \pm 6.3\%$ of baseline; n=4; p<0.05; Fig 3B). Similarly, treatment with 25 nM leptin (116-130) before A β_{1-42} resulted in a robust increase in synaptic transmission (to $136 \pm 5.5\%$ of baseline; n=6; p<0.05; Fig 3C). In contrast, HFS failed to increase synaptic efficacy ($107 \pm 3.3\%$ of baseline; n=5; p>0.05) in slices exposed to A β_{1-42} and leptin (22-56; Fig 4D), although some STP was observed. Thus these data indicate that like leptin, leptin (116-130) prevents the detrimental effects of A β_{1-42} on hippocampal synaptic plasticity.

Leptin (116-130) reverses A β_{1-42} -induced LTD.

It is known that oligomeric A β promotes the induction of LTD (Shankar et al, 2008) and that exposure to a low concentration of leptin (10 nM) prevents facilitation of hippocampal LTD by A β_{1-42} (Doherty et al, 2013). Thus we assessed if either of the leptin fragments mirror leptin action. Initially we verified that leptin prevented A β_{1-42} -induced LTD. In agreement

with previous studies (Doherty et al, 2013), application of the sub-threshold LFS paradigm failed to induce LTD in vehicle-treated slices ($94 \pm 5.6\%$ of baseline; $n=5$; $p>0.05$), whereas robust LTD ($73 \pm 3.8\%$ of baseline; $n=5$; $p<0.001$) was induced in $A\beta_{1-42}$ -treated slices (Fig 4A). Furthermore, leptin (10nM) reduced the magnitude of $A\beta_{1-42}$ -induced LTD such that LFS depressed synaptic transmission to $101 \pm 5.3\%$ of baseline in leptin-treated slices ($n=4$; $p>0.05$; Fig 4B). To establish if the leptin fragments mirrored leptin action the effects of leptin (116-130) or leptin (22-56) were also examined. Application of 10nM leptin (116-130) or leptin (22-56) had no significant effect on basal synaptic transmission ($n=6$ for leptin (116-130) and $n=5$ for leptin (22-56)). Application of 10nM leptin (22-56) failed to alter the magnitude of $A\beta_{1-42}$ -induced LTD such that synaptic transmission was depressed to $73 \pm 4.0\%$ of baseline ($n=5$; $p<0.001$; Fig 4D). In contrast, a significant reduction in the magnitude of $A\beta_{1-42}$ -induced LTD ($94 \pm 5.2\%$ of baseline; $n=5$; $p>0.05$; Fig 4C) was observed in hippocampal slices treated with leptin (116-130; 10nM; 45 min) indicating that leptin (116-130) also inhibits facilitation of LTD by $A\beta_{1-42}$. Thus these data indicate that leptin (116-130), but not leptin (22-56) reverses $A\beta_{1-42}$ -induced facilitation of LTD.

Leptin (116-130) prevents $A\beta$ -induced internalization of GluA1

Previous studies indicate that $A\beta$ promotes internalisation of the AMPA receptor subunit, GluA1 (Hsieh et al, 2006; Liu et al, 2010); an effect that is prevented by leptin (Doherty et al, 2013). To determine if the leptin fragments mirror this effect, the cell surface density of GluA1 was probed in cultured hippocampal neurons (Moult et al, 2010). In accordance with previous studies (Doherty et al, 2013), treatment with $A\beta_{1-42}$ (500 nM; 20 min) significantly attenuated (to $70 \pm 2\%$ of control) GluA1 surface expression compared with control ($A\beta_{42-1}$ -treated) hippocampal neurons ($n=48$; $p<0.001$; Fig. 4E). Application of a low concentration of leptin (10nM) induced a small increase in GluA1 surface expression ($113 \pm 7\%$ of control; $n=36$; $p<0.01$). However, in leptin-treated neurons, $A\beta_{1-42}$ failed to significantly alter GluA1

surface expression ($101 \pm 4\%$ of control; $n=36$; $P>0.05$). In contrast, treatment with a low concentration (10nM) of leptin (22-56) had no significant effect on GluA1 trafficking *per se* ($n=36$; $p>0.05$). Moreover, leptin (22-56) failed to prevent the effects of A β on GluA1 trafficking as surface GluA1 was reduced to $51 \pm 2\%$ of control ($n=36$; $p<0.001$) in the presence of A β and leptin (22-56; Fig 4E). Conversely, after exposure to leptin (116-130), A β_{1-42} failed to significantly reduce GluA1 surface expression ($97 \pm 4\%$ of control; $n=36$; $p>0.05$). These data indicate that leptin (116-130) mirrors leptin action by preventing A β_{1-42} -induced internalisation of GluA1 in cultured hippocampal neurons.

Leptin (116-130) prevents copper and A β -induced cell death

We have demonstrated previously that leptin attenuates cortical neuronal death triggered by A β_{1-42} or divalent copper ions (Doherty et al, 2013). To determine whether leptin (116-130) has neuro-protective actions, the effects of leptin (116-130) on the viability of differentiated human neural cells (SH-SY5Y) was examined after exposure to either 5 μ M CuCl₂ or 10 μ M A β_{1-42} . Cells were treated with the toxin alone or with a range of concentrations (10 - 0.1nM) of leptin or leptin (116-130). Determination of membrane leakage by lactate dehydrogenase (LDH) assay revealed a significant reduction in LDH release after treatment with either leptin or leptin (116-130; both 0.1-10nM). Thus for CuCl₂-treated cells, 10nM leptin reduced LDH release by $39.5 \pm 2.73\%$ compared to CuCl₂ alone ($n=5$; $p<0.001$; Fig 5A); an effect that was mirrored by 10nM leptin (116-130; $45.6 \pm 2.92\%$ ($n=5$; $p<0.001$; Fig 5A). In A β_{1-42} treated cells, leptin also significantly reduced LDH release by $31.8 \pm 13.2\%$ (0.1nM) and $47.9 \pm 7.45\%$ (10nM), respectively ($n=5$; $p<0.001$; Fig 5B). Similarly, application of 0.1nM or 10nM leptin (116-13) decreased LDH release by $26.7 \pm 17.3\%$ or $46.6 \pm 9\%$, respectively ($n=5$; $p<0.05$; Fig 5B). Thus like leptin, treatment with leptin (116-130) reduces neuronal death in response to AD-linked toxins *in vitro*.

In parallel studies a crystal violet assay was used to verify these findings by assessing cell number. In CuCl₂-treated cells, there was a concentration-dependent increase in the survival of cells treated with either leptin or leptin (116-130). Thus, treatment with leptin (0.1nM) resulted in a $16.7 \pm 3.4\%$ increase in cell number and this increased to $43.4 \pm 9.2\%$ in the presence of 10nM leptin (n=5; p<0.01). Similarly, exposure to 0.1nM or 10nM leptin (116-130) increased cell number by $27.8 \pm 10.6\%$ and $39.9 \pm 13.5\%$, respectively (n=5; Fig 5C). Treatment with leptin (116-130) also mirrored leptin action by increasing cell viability in A β ₁₋₄₂-treated cultures as cell number increased by $19.2 \pm 15\%$ and $44.3 \pm 7.5\%$ after treatment with 0.1nM or 10nM leptin, respectively (n=5 for each; p<0.01). Exposure to leptin (116-130) also resulted in significant increases in cell number (0.1nM: $29.4 \pm 9.4\%$ increase; 10nM: $51.8 \pm 6.3\%$ increase; n=5 for both; p<0.001; Fig 5D).

As leptin (22-56) has biological activity in other systems, the specificity of the leptin (116-130) fragment in promoting cell survival was examined by determining whether leptin (22-56) inhibited neuronal death induced by A β ₁₋₄₂. In contrast to leptin (116-130), treatment with leptin (22-56) had no effect on the viability of cells exposed to A β ₁₋₄₂ ($41.2 \pm 5.8\%$ survival following A β ₁₋₄₂ treatment and $48.6 \pm 9.3\%$ in A β ₁₋₄₂ with 10nM leptin (22-56) treated cells; n=5; p>0.5; data not shown). These data reveal a potent neuroprotective effect of the leptin fragment (116-130) that is comparable to the survival actions of leptin. Moreover, this anti-apoptotic response is specific to leptin (116-130) as leptin (22-56) failed to influence neuronal viability.

The neuroprotective effects of leptin (116-130) involve activation of STAT3 and PI3-kinase-dependent signalling pathways

Our previous studies indicate a crucial role for STAT3 and PI3-kinase/Akt signalling in the neuroprotective actions of leptin (Doherty et al, 2013). To determine whether leptin (116-

130) acts via similar signalling cascades we examined the effects of pharmacological inhibitors of STAT3 (WP1066) or PI3-kinase (wortmannin). In A β_{1-42} -treated SH-SY5Y cells, application of either inhibitor significantly reduced the ability of leptin (116-130) to alleviate neuronal death. When neurons were treated with the STAT3 inhibitor, an $18.3 \pm 3.2\%$ increase in LDH release in leptin (116-130) and A β_{1-42} -treated cultures was observed, which is similar to the $26.7 \pm 4.4\%$ increase observed with A β_{1-42} alone ($n=5$; $p>0.5$; Fig 6A). Thus STAT3 inhibition blocks the neuroprotective actions of leptin (116-130), suggesting a role for STAT3 in this process. Furthermore, following inhibition of PI3-kinase with wortmannin an $26.9 \pm 9.8\%$ increase in LDH release was observed in leptin (116-130) and A β_{1-42} -treated cells which is not significantly different from cells treated with A β_{1-42} alone ($n=5$; $p>0.5$; Fig 6B). Thus these data also indicate a role for PI3-kinase in mediating the neuroprotective actions of leptin (116-130).

To verify that leptin (116-130) directly activates these signalling pathways, SH-SY5Y cells were exposed to 1nM leptin (116-130; 3 h) or left untreated prior to protein extraction for ELISA. The ratio of phosphorylated STAT3 to pan STAT3 increased markedly following leptin (116-130) administration ($n=3$; $p<0.01$; Fig 6C). Similarly an increase in the ratio of phosphorylated Akt to pan Akt was observed following exposure to leptin (116-130; $n=3$; $p<0.01$; Fig 6D). These data indicate that leptin (116-130) reduces cell death by a mechanism involving activation of STAT3 and PI3-kinase. Furthermore, exposure to leptin (116-130) resulted in a significant increase in the active components of these signalling cascades.

Leptin (116-130) enhances episodic-like memory

The current data demonstrate that leptin (116-130) enhances hippocampal synaptic plasticity mechanisms and has neuroprotective effects. To further assess its therapeutic potential we next asked if this fragment has similar cognitive enhancing properties to the whole leptin

molecule. Previous studies indicate that leptin enhances hippocampal-dependent memory (Oomura et al, 2006; Farr et al, 2006), whereas resistance to leptin results in impaired spatial memory (Li et al, 2002). We used the object-place-context (OPC) recognition task which models human episodic memory, the first cognitive process to be compromised in the early stages of AD (Swainson et al, 2001). Performance on this task has been shown to be impaired in murine models of AD (Davis et al, 2013) and is compromised in animals with lesions of hippocampus (Langston and Wood, 2010) and lateral entorhinal cortex (Wilson et al, 2013). The task is based on the object recognition paradigm and models the integrated aspect of human episodic memory by exposing rodents to novel combinations of objects, the spatial locations in which they are experienced and the contextual features of the environment (see Fig 7A; Eacott and Norman, 2004). 42 C57/BL6 mice were habituated to a testing environment and then trained on object recognition, object-place recognition and object-context recognition. Following training mice were tested on 4 days of the episodic-like OPC task. On these days mice were assigned to one of three groups (control, leptin, or fragment) and on each day mice were given 100 μ l IP injections of saline, 7.8nM/ml leptin, or 7.8nM/ml leptin (116-130) 30 min prior to testing. One-way ANOVA on the discrimination indices during the first minute of the test phase revealed a significant effect of group (Fig 7B; $F_{(2,41)} = 4.318$, $p=0.02$). Posthoc comparisons (Tukey's HSD) revealed that both the leptin and leptin (116-130) treated mice showed enhanced performance on the task relative the control group ($p<0.05$) and did not differ from each other. One sample t-tests confirmed that all groups performed significantly better than chance level performance ($p<0.01$). Finally analysis of the overall exploration time in both sample and test phases of the task revealed no change in total levels of exploration between groups demonstrating that neither leptin nor leptin (116-130) produced a non-specific change in exploration behaviour (Fig 7C; $p>0.05$). The increased discrimination index was driven by an increase in exploration of the novel

object combined with a decrease in the exploration of the familiar object (Figure 7D).

Together these data indicate that leptin (116-130) mirrors leptin's action by enhancing performance in episodic-like memory tasks.

Discussion

It is well established that the hormone leptin circulates in the plasma and enters the brain via transport across the blood brain barrier. In the hypothalamus, leptin plays a major role in regulating food intake and body weight (Spiegelman and Flier, 2001). However, the central actions of the hormone leptin are not restricted to the hypothalamus and the regulation of energy homeostasis. Indeed, a number of extra-hypothalamic brain regions, including the hippocampus display high levels of leptin receptor expression (Irving and Harvey 2014).

Leptin mRNA and protein are also highly expressed in the hippocampal formation (Morash et al, 1999) and emerging evidence suggests brain-specific production of leptin (Eikelis et al, 2006). Thus it is likely that a combination of locally released leptin as well as peripherally-derived leptin reach hippocampal synapses and can influence synaptic function. Indeed, numerous studies indicate that leptin has potential cognitive enhancing properties as it readily facilitates the cellular events underlying hippocampal learning and memory. Thus, leptin has rapid effects on activity-dependent synaptic plasticity, glutamate receptor trafficking and dendritic morphology (Irving and Harvey, 2014). In addition, several studies have identified neuroprotective effects of leptin as the viability of central and peripheral neurons is markedly influenced by this hormone (Doherty et al, 2008; Weng et al, 2007; Guo et al, 2008). Recent clinical evidence has established a link between circulating leptin levels and the incidence of AD (Power et al, 2001; Lieb et al, 2009), which has fuelled the possibility of using the leptin system as a novel therapeutic target in AD. Indeed, treatment of various AD models with leptin prevents the detrimental effects of A β that occur at both early and late stages of the

disease (Fewlass et al, 2004; Doherty et al, 2013; Farr et al, 2006). However, as leptin is a very large peptide, developing small leptin-like molecules may be a better therapeutic approach. Several fragments of the leptin peptide are biologically active and mirror the anti-obesity effects of leptin (Grasso et al, 1997; Grasso et al, 2001; Rozhavskaya-Arena et al, 2000). However, the cognitive enhancing and neuroprotective effects of the leptin fragments are not known. Here we provide the first compelling evidence that leptin (116-130), but not leptin (22-56), has a potent effect on hippocampal synaptic function as it promotes trafficking of AMPA receptors to synapses and facilitates hippocampal synaptic plasticity. Moreover in cellular models that mimic amyloid toxicity, leptin (116-130), but not leptin (22-56) prevents the aberrant effects of A β on hippocampal synaptic function and neuronal viability. These findings indicate that one particular leptin fragment, namely (116-130), mirrors the beneficial actions of leptin in preventing the detrimental effects of A β at the early and late stages of AD. Finally we have shown that the leptin fragment that enhances hippocampal synaptic plasticity and has neuroprotective effects, namely leptin (116-130), is also a cognitive enhancer as it improves performance on tests of episodic memory.

Here we show that, in accordance with previous studies (Wayner et al, 2004; Shanley et al, 2001), NMDA receptor-dependent synaptic plasticity is enhanced by leptin as treatment with leptin promoted conversion of STP into a persistent increase in synaptic transmission in juvenile hippocampal slices. Similarly exposure to the leptin fragment (116-130) readily facilitated synaptic plasticity as an increase in synaptic strength was evident in leptin (116-130), but not leptin (22-56)-treated slices. We have shown that the efficacy of excitatory synaptic transmission is also regulated by leptin in adult hippocampus (Moult et al, 2010). In agreement with this, application of either leptin or leptin (116-130) to adult hippocampal slices resulted in the induction of a persistent increase in synaptic transmission. In contrast, however leptin (22-56) failed to alter excitatory synaptic strength in adult hippocampus.

AMPA receptor trafficking is pivotal for activity-dependent synaptic plasticity (Collingridge et al, 2004) and leptin regulates trafficking of GluA1 to synapses (Moult et al, 2010). In this study, treatment with either leptin or leptin (116-130) increased GluA1 surface expression in cultured hippocampal neurons, whereas leptin (22-56) was without effect. In co-localisation studies, the density of GluA1 subunits associated with synapses was increased after application of leptin or leptin (116-130), suggesting that leptin (116-130) parallels the actions of leptin by boosting the synaptic insertion of AMPA receptors. We have shown that leptin-driven trafficking of GluA1 involves inhibition of PTEN (Moult et al, 2010). Similarly in this study, the ability of leptin (116-130) to influence GluA1 trafficking involves inhibition of PTEN, as application of the PTEN inhibitor bpV blocked the increase in GluA1 surface expression induced by leptin (116-130) in hippocampal neurons. These data indicate that like leptin, treatment with leptin (116-130) promotes GluA1 trafficking to hippocampal synapses via inhibition of PTEN. Thus, overall these data indicate that the leptin fragment (116-130) mirrors the actions of leptin as it markedly influences the cellular events underlying learning and memory by regulating AMPA receptor trafficking.

It is known that A β inhibits the induction of hippocampal LTP (Shankar et al, 2008), and this detrimental effect of A β_{1-42} is reversed by leptin (Doherty et al, 2013). Similarly, leptin (116-130) reversed the acute effects of A β_{1-42} in this study as synaptic plasticity was readily induced in hippocampal slices exposed to leptin (116-130) and A β_{1-42} . Contrastingly, application of leptin (22-56) failed to prevent the detrimental effects of A β_{1-42} as no increase in synaptic strength was induced after exposure to A β_{1-42} and leptin (22-56). However, in slices exposed to leptin (22-56) post-tetanic potentiation (PTP) and some STP was observed after HFS, suggesting that this fragment may influence the transient enhancement of synaptic strength induced by HFS. As PTP and STP are thought to involve presynaptic expression

mechanisms (Zucker and Regehr, 2002; Lauri et al, 2007), it is feasible that leptin (22-56) can act pre-synaptically to influence glutamate release mechanisms.

Several studies indicate that A β ₁₋₄₂ also facilitates the induction of hippocampal LTD (Shankar et al, 2008; Li et al, 2009), and this effect is also reversed by leptin (Doherty et al, 2013). In accordance with these findings, treatment with leptin reduced the magnitude of LTD in A β ₁₋₄₂-treated slices. Similarly, leptin (116-130), but not leptin (22-56) attenuated the effects of A β ₁₋₄₂ as the magnitude of LTD was significantly decreased in the presence of leptin (116-130). Moreover, application of either leptin or leptin (116-130) inhibited A β ₁₋₄₂-driven AMPA receptor removal from hippocampal synapses, whereas treatment with leptin (22-56) was without effect. Thus overall these data demonstrate that leptin (116-130) mirrors the actions of leptin in counteracting the detrimental acute effects of A β ₁₋₄₂ on hippocampal synaptic function.

Evidence is growing that leptin has neuroprotective actions in various models of neurodegenerative disease. In Parkinson's disease models, treatment with leptin protects dopaminergic neurons from various toxic insults (Doherty et al, 2008; Weng et al, 2007), whereas in AD models of amyloid toxicity, leptin increases neuronal viability via activation of STAT3 and PI3-kinase signalling (Doherty et al, 2013; Doherty et al, 2008; Guo et al, 2008). In this study, leptin and leptin (116-130) enhanced the survival of human neural (SH-SY5Y) cells treated with either A β ₁₋₄₂ or Cu²⁺. Conversely no change in cell viability was evident after treatment with leptin (22-56), thereby providing further evidence that leptin (116-130) but not leptin (22-56) mirrors the protective actions of leptin.

In these studies, we reveal that signalling via PI3-kinase and STAT3 is essential for leptin (116-130)-mediated neuroprotection as selective inhibition of these pathways eliminated the protective effects of leptin (116-130). Moreover, direct activation of key components of PI3-

kinase and STAT3 signalling pathways was observed following administration of leptin (116-130). As stimulation of both PI3-kinase (Doherty et al, 2013; Doherty et al, 2008) and STAT3 signalling cascades (Doherty et al, 2013; Guo et al, 2008) mediate the neuroprotective actions of leptin, these data indicate that leptin (116-130) is activating the same signalling pathways as the full length leptin peptide to induce neuronal survival. This provides further evidence that leptin (116-130) is mirroring the neuronal effects of leptin.

These studies demonstrate that the 116-130 fragment of the leptin molecule enhances hippocampal synaptic plasticity and has neuroprotective effects. As such this fragment is a very interesting therapeutic target to treat memory dysfunction and protect against neurodegeneration in the early stages of AD. To test the functional implications of the effects of leptin 116-130 we examined the effects of acute doses of this fragment on a test of episodic-like memory. This test is particularly appropriate as it models the type of memory that is first compromised in AD. Performance on the task has been shown to be impaired in rodents with damage to the lateral entorhinal cortex (Wilson et al, 2013), the first region to be damaged in AD, and the hippocampus (Langston and Woods, 2010). It has also been shown that the triple transgenic murine model of AD show impaired performance on this task at 6 months of age (Davis et al, 2013). The current data demonstrate powerful cognitive enhancing effects of both leptin and leptin (116-130) as both groups performed significantly better than controls on the OPC task. This is the first time that leptin has been shown to enhance the specific type of memory that degrades in AD and the fact that this cognitive enhancement is also produced by leptin (116-130) suggests that this fragment is a viable tool to treat memory dysfunction caused by damage to the hippocampal-entorhinal network. Recent studies indicate that administration of leptin also protects against A β -induced impairments in spatial memory tasks (Tong et al, 2015). Thus it is feasible that administration

of leptin (116-130) will also mirror the effects of leptin and protect against the chronic effects of A β on hippocampal-dependent learning and memory.

The current experiments demonstrated enhancement of memory for object-place-context associations. Enhancement of this hippocampal dependent task is consistent with our findings showing enhancement of hippocampal synaptic plasticity but it remains a possibility that leptin 116-130 may also enhance simpler forms of recognition memory such as object recognition or object-place recognition. These simpler forms of recognition memory are dependent on other areas of the medial temporal lobe network and so future work could examine whether the cognitive enhancement is specific to the hippocampus or also affects the surrounding cortical inputs. One other consideration is the anxiolytic properties of leptin that have been reported in both normal (Liu et al. 2010) and chronically stressed rats (Lu et al. 2006). Reduced anxiety could potentially affect performance on the spontaneous recognition tasks as less anxious animals may explore more freely. This was not found to be the case in the current study as the levels of exploration in both sample and test phases of the OPC experiment were not different between groups. This is not surprising as animals had extensive handling and pre-training before the OPC test and so levels of anxiety would have been very low in all animals.

In conclusion, these data indicate that the leptin (116-130) fragment mirrors the cognitive enhancing effects of leptin as it promotes trafficking of the AMPA receptor subunit GluA1 to synapses, facilitates hippocampal synaptic plasticity and improves performance in an episodic-like memory task. In addition, leptin (116-130) counteracts the detrimental effects of A β ₁₋₄₂ on hippocampal synaptic function and neuronal viability in various cellular models of amyloid toxicity. Our findings not only reinforce the consensus that the leptin system is an important therapeutic target in AD, but also establish that leptin (116-130) may be useful in the development of leptin-mimetic agents for therapeutic use.

Funding

J.H. is funded by The Anonymous Trust and Cunningham Trust. GD is funded by ARUK, DR received a University of St Andrews Research Internship. JAA is funded by the Carnegie Trust.

Notes.

We would like to thank Matthew Buchan and Xiaoyue Zhu for help with behavioural testing.

References.

Anderson WW, Collingridge GL. 2007. Capabilities of the WinLTP data acquisition program extending beyond basic LTP experimental functions. *J Neurosci Meth.* 162:346–356.

Bliss TV, Collingridge GL. 1993. A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 361:31-9.

Collingridge GL, Isaac JT, Wang YT. 2004. Receptor trafficking and synaptic plasticity. *Nat Rev Neurosci.* 5:952-62.

Davis KE, Eacott MJ, Easton A, Gigg J. 2013. Episodic-like memory is sensitive to both Alzheimer's-like pathological accumulation and normal ageing processes in mice. *Behav Brain Res.* 254:73-82.

Doherty GH, Beccano-Kelly D, Yan SD, Gunn-Moore FJ, Harvey J. 2013. Leptin prevents hippocampal synaptic disruption and neuronal cell death induced by amyloid β . *Neurobiol Aging* 34:226-37.

Doherty GH, Oldreive C, Harvey J. 2008. Neuroprotective actions of leptin on central and peripheral neurons in vitro. *Neurosci.* 154:1297-307

Eacott MJ, Norman G. 2004. Integrated memory for object, place, and context in rats: a possible model of episodic-like memory? *J Neurosci.* 24:1948-53.

Eikelis N, Wiesner G, Lambert G, Esler M. 2006. Brain leptin resistance in human obesity revisited. *Regul Pept.* 139:45-51.

Farr SA, Banks WA, Morley JE. 2006. Effects of leptin on memory processing. *Peptides* 27: 1420-5.

Fewlass DC, Noboa K, Pi-Sunyer FX, Johnston JM, Yan SD, Tezapsidis N. 2004. Obesity-related leptin regulates Alzheimer's Abeta. *FASEB J.* 18:1870-1878.

Grasso P, Leinung MC, Ingher SP, Lee DW. 1997. In vivo effects of leptin-related synthetic peptides on body weight and food intake in female ob/ob mice: localization of leptin activity to domains between amino acid residues 106-140. *Endocrinol.* 138:1413-8.

Grasso P, Rozhavskaya-Arena M, Leinung MC, Lee DW. 2001. [D-LEU-4]-OB3, a synthetic leptin agonist, improves hyperglycemic control in C57BL/6J ob/ob mice. *Regul Pept.* 101:123-9.

Guo Z, Jiang H, Xu X, Duan W, Mattson MP. 2008. Leptin-mediated cell survival signaling in hippocampal neurons mediated by JAK STAT3 and mitochondrial stabilization. *J Biol Chem.* 283:1754-63.

Hsieh H, Boehm J, Sato C, Iwatsubo T, Tomita T, Sisodia S, Malinow R. 2006. AMPAR removal underlies Abeta-induced synaptic depression and dendritic spine loss. *Neuron*. 52:831-843.

Irving AJ, Harvey J. 2014. Leptin regulation of hippocampal synaptic function in health and disease. *Phil Trans R Soc. B* 369:20130155.

Langston RF, Wood ER. 2010. Associative recognition and the hippocampus: differential effects of hippocampal lesions on object-place, object-context and object-place-context memory. *Hippocampus*. 20:1139-53

Lauri SE, Palmer M, Segerstrale M, Vesikansa A, Taira T, Collingridge GL. 2007. Presynaptic mechanisms involved in the expression of STP and LTP at CA1 synapses in the hippocampus. *Neuropharmacol*. 52:1-11

Li XL, Aou S, Oomura Y, Hori N, Fukunaga K, Hori T. 2002. Impairment of long-term potentiation and spatial memory in leptin receptor-deficient rodents. *Neurosci*. 113:607-15.

Li S, Hong S, Shepardson NE, Walsh DM, Shankar GM, Selkoe D. 2009. Soluble oligomers of amyloid Beta protein facilitate hippocampal long-term depression by disrupting neuronal glutamate uptake. *Neuron*. 62:788-801.

Lieb W, Beiser AS, Vasan RS, Tan ZS, Au R, Harris TB, Roubenoff R, Auerbach S, DeCarli C, Wolf PA, Seshadri S 2009. Association of plasma leptin levels with incident Alzheimer disease and MRI measures of brain aging. *JAMA*. 302:2565-2572.

Liu J1, Garza JC, Bronner J, Kim CS, Zhang W, Lu XY. 2010. Acute administration of leptin produces anxiolytic-like effects: a comparison with fluoxetine. *Psychopharmacol.* 207(4):535-45.

Liu SJ, Gasperini R, Foa L, Small DH. 2010. Amyloid-beta decreases cell-surface AMPA receptors by increasing intracellular calcium and phosphorylation of GluR2. *J Alzheimers Dis.* 21:655-666.

Luo X, McGregor G, Irving AJ, Harvey J. 2015. Leptin Induces a Novel Form of NMDA Receptor-Dependent LTP at Hippocampal Temporoammonic-CA1 Synapses. *eNeuro* 2(3):1-7.

Lu XY, Kim CS, Frazer A, Zhang W. 2006. Leptin: a potential novel antidepressant. *Proc Natl Acad Sci U S A.* 103:1593-1598.

Morash B, Li A, Murphy PR, Wilkinson M, Ur E. 1999. Leptin gene expression in the brain and pituitary gland. *Endocrinol.* 140:5995-8.

Moult PR, Cross A, Santos SD, Carvalho AL, Lindsay Y, Connolly CN, Irving AJ, Leslie NR, Harvey J. 2010. Leptin regulates AMPA receptor trafficking via PTEN inhibition. *J Neurosci.* 30:4088-101.

Moult PR, Harvey J. 2011. NMDA receptor subunit composition determines the polarity of leptin-induced synaptic plasticity. *Neuropharmacol.* 61:924-36.

Oldreive CE, Doherty GH. 2010. Effects of tumour necrosis factor-alpha on developing cerebellar granule and Purkinje neurons in vitro. *J Mol Neurosci.* 42:44-52.

O'Malley D, MacDonald N, Mizielinska S, Connolly CN, Irving AJ, Harvey J. 2007. Leptin promotes rapid dynamic changes in hippocampal dendritic morphology. *Mol Cell Neurosci.* 35:559-72

Oomura Y, Hori N, Shiraishi T, Fukunaga K, Takeda H, Tsuji M, Matsumiya T, Ishibashi M, Aou S, Li XL, Kohno D, Uramura K, Sougawa H, Yada T, Wayner MJ, Sasaki K. 2006. Leptin facilitates learning and memory performance and enhances hippocampal CA1 long-term potentiation and CaMK II phosphorylation in rats. *Peptides.* 27:2738-49

Power DA, Noel J, Collins R, O'Neill D. 2001. Circulating leptin levels and weight loss in Alzheimer's disease patients. *Dement Geriatr Cogn Disord.* 12:167-70.

Rose GM, Dunwiddie TV. 1986. Induction of hippocampal long-term potentiation using physiologically patterned stimulation. *Neurosci Lett.* 69:244-8.

Rozhavskaya-Arena M, Lee DW, Leinung MC, Grasso P. 2000. Design of a synthetic leptin agonist: effects on energy balance, glucose homeostasis, and thermoregulation. *Endocrinol.* 141:2501-7.

Samson WK, Murphy TC, Robison D, Vargas T, Tau E, Chang JK. 1996. A 35 amino acid fragment of leptin inhibits feeding in the rat. *Endocrinol.* 137:5182-5.

Schmid AC, Byrne RD, Vilar R, Woscholski R. 2004. Bisperoxovanadium compounds are potent PTEN inhibitors. *FEBS Lett.* 566:35-8.

Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, Smith I, Brett FM, Farrell MA, Rowan MJ, Lemere CA, Regan CM, Walsh DM, Sabatini DL, Selkoe DJ. 2008.

Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med.* 14:837-842.

Shanley LJ, Irving AJ, Harvey J. 2001. Leptin enhances NMDA receptor function and modulates hippocampal synaptic plasticity. *J Neurosci.* 21:RC186.

Simmons LK, May, PC, Tomaselli, KJ, Rydel RE, Fuson KS, Brigham EF, Wright S, Lieberburg I, Becker GW, Brems DN, Li WY 1994. Secondary structure of amyloid beta peptide correlates with neurotoxic activity in vitro. *Mol Pharmacol*, 45:373–379

Spiegelman BM, Flier JS. 2001. Obesity and the regulation of energy balance. *Cell.* 104:531–43.

Stranahan AM, Mattson MP 2012. Metabolic reserve as a determinant of cognitive aging. *J Alzheimers Dis.* 30:S5-13.

Swainson R, Hodges JR, Galton CJ, Semple J, Michael A, Dunn BD, Iddon JL, Robbins TW Sahakian BJ. 2001. Early detection and differential diagnosis of Alzheimer's disease and depression with neuropsychological tasks. *Dement Geriat Cogn Disord.* 12:265-280.

Tong JQ, Zhang J, Hao M, Yang J, Han YF, Liu XJ, Shi H, Wu MN, Liu QS, Qi JS. 2015. Leptin attenuates the detrimental effects of β -amyloid on spatial memory and hippocampal later-phase long term potentiation in rats. *Horm Behav.* 73:125-30.

Walsh DM, Klyubin I, Fadeeva JV, Cullen WK, Anwyl R, Wolfe MS, Rowan MJ, Selkoe DJ 2002. Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo. *Nature.* 416:535-9.

Wayner MJ, Armstrong DL, Phelix CF, Oomura Y. 2004. Orexin-A (Hypocretin-1) and leptin enhance LTP in the dentate gyrus of rats in vivo. *Peptides*. 25:991-6.

Weng Z, Signore AP, Gao Y, Wang S, Zhang F, Hastings T, Yin XM, Chen J. 2007. Leptin protects against 6-hydroxydopamine-induced dopaminergic cell death via mitogen-activated protein kinase signaling. *J Biol Chem*. 282:34479-91.

Wilson DI, Watanabe S, Milner H, Ainge JA. 2013. Lateral entorhinal cortex is necessary for associative but not non associative recognition memory. *Hippocampus*. 23:1280-90

Zucker RS, Regehr WG. 2002. Short-term synaptic plasticity. *Annu Rev Physiol*. 64:355-405.

Figure Legends

Figure 1.

Leptin (116-130) promotes conversion of STP into a persistent increase in synaptic transmission in juvenile hippocampus.

A. Pooled data showing that primed burst stimulation (indicated by the arrow) delivered in the absence of leptin induced STP (filled circle) in juvenile hippocampal slices. In contrast, in leptin-treated slices (open circle) the same stimulation paradigm resulted in a persistent increase in synaptic transmission. In this and subsequent figures, each point is the average of four successive responses. Top, Representative synaptic records (average of four consecutive records) are shown for the times indicated. B, During exposure to leptin (116-130; 50nM), application of the primed burst stimulation paradigm resulted in a persistent increase in

synaptic transmission (B), whereas only STP was evident in slices treated with leptin (22-56; C).

Figure 2.

In adult, leptin (116-130) induces a persistent increase in synaptic efficacy and increases GluA1 trafficking to hippocampal synapses. .

A. Pooled data showing that application of leptin (116-130) results in a persistent increase in excitatory synaptic transmission in adult hippocampal slices. B, In contrast application of leptin (22-56) failed to alter excitatory synaptic strength. C, Representative confocal images of surface GluR1 staining in control cultured hippocampal neurons and after exposure to leptin, leptin (116-130) or leptin (22-56). Leptin (116-130) mirrors the effects of leptin by increasing GluA1 surface labelling. Scale bars, 10 μ m. D, Pooled data showing relative changes in surface GluA1 labelling in control conditions, and after exposure to leptin, leptin (116-130) or leptin (22-56) in hippocampal neurons. E, Pooled data of the percent co-localisation of surface GluA1 and synaptophysin immunolabelling in cultured hippocampal neurons. Leptin (116-130), but not leptin (22-56) increased GluA1 surface labelling associated with synapses. F. Pooled data of relative changes in surface GluA1 labelling in hippocampal cultures in control conditions, after bpV, leptin or leptin (116-130) treatment, and in the presence of bpV and either leptin, or leptin (116-130). Inhibition of PTEN mimicked and occluded the effects of leptin (116-130).

Figure 3.

Leptin (116-130) inhibits the aberrant effects of amyloid- β (A β)₁₋₄₂ on hippocampal synaptic plasticity in juvenile slices.

A, Pooled data showing that HFS (indicated by the arrow) induces synaptic plasticity in $A\beta_{42-1}$ -treated (open circles) slices, whereas $A\beta_{1-42}$ inhibits synaptic plasticity (filled circles). B, Exposure to leptin prevented $A\beta_{1-42}$ -inhibition of synaptic plasticity. Treatment with leptin (116-130; C), but not leptin (22-56; D) reversed $A\beta_{1-42}$ -inhibition of synaptic plasticity.

Figure 4.

Leptin (116-130) prevents $A\beta$ -induced AMPA receptor internalisation and facilitation of hippocampal LTD.

A, Pooled data showing that the sub-threshold low frequency stimulation (indicated by the arrow) failed to induce long-term depression (LTD) in $A\beta_{42-1}$ -treated (open circles) slices, whereas robust LTD (filled circles) is induced in $A\beta_{1-42}$ -treated juvenile slices. B, Exposure to leptin prevented $A\beta_{1-42}$ -induced LTD. Similarly treatment with leptin (116-130; C), but not leptin (22-56; D) prevented $A\beta_{1-42}$ -induced LTD. E, Pooled data showing relative changes in GluA1 surface labelling in cultured hippocampal neurons in control ($A\beta_{42-1}$) conditions and after treatment with leptin, $A\beta_{1-42}$, leptin (116-130), leptin (22-56) and in the presence of $A\beta_{1-42}$ plus either leptin, leptin (116-130) or leptin (22-56), respectively. Leptin and leptin (116-130) prevent $A\beta$ -driven internalisation of GluA1.

Figure 5.

Leptin (116-130) prevents copper and $A\beta$ -induced cell death in human SH-SY5Y cells.

A, Pooled data revealing that leptin and leptin (116-130) prevent LDH release induced by administration of $5\mu\text{M}$ CuCl_2 . B, Similar pooled data was obtained for cultures treated with $10\mu\text{M}$ $A\beta_{1-42}$. C, Pooled data showing that in $5\mu\text{M}$ CuCl_2 treated cultures enhanced numbers of cells are detected with a crystal violet assay when cultures are co-treated with either leptin

or leptin (116-130) with a similar trend observed when cultures were induced to die with 10 μ M A β ₁₋₄₂, D.

Figure 6.

The neuroprotective effects of leptin (116-130) involve activation of STAT3 and PI3-kinase-dependent signalling pathways

A, Pooled data revealing that treatment of SH-SY5Y cells with the STAT-3 inhibitor WP1066 (5 μ M) prevented leptin (116-130)-mediated neuroprotection from the effects of 10 μ M A β ₁₋₄₂. B, Pooled data from LDH assays demonstrating that the PI3-kinase inhibitor wortmannin (50nM) attenuated leptin (116-130)-mediated neuroprotection against 10 μ M A β ₁₋₄₂. C, Pooled data from ELISA assays showing that leptin (116-130) stimulates phosphorylation of STAT3. D, Pooled data from ELISA assays showing that leptin (116-130) stimulates phosphorylation of Akt.

Figure 7.

Leptin (116-130) enhances episodic-like memory.

A. Object-place-context task used to assess episodic-like memory. There are 2 sample phases in which mice are exposed to different copies of two different objects (star and hexagon) and allowed to explore for 3 mins. In the test phase they see two new copies of one of the objects. The arrow points to the object that has not been previously seen in that place within that context. B. Mean \pm SEM discrimination index for the three groups. * indicates $p < 0.05$. C. Total exploration time in the test phase is not different between groups. D. Exploration of the novel and familiar objects in the test phase.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

